



NATIONAL
MARINE
SANCTUARY
FOUNDATION



CAL POLY
SAN LUIS OBISPO




Carbon and Nitrogen Changes due to Soil Warming



Dana Coppernoll-Houston¹, Chris Potter²

¹Federal Way High School

²NASA Ames Research Center



Background: Soil


- Soil is a complex ecosystem
 - Minerals
 - Water
 - Gasses
 - Organic Matter
 - Microorganisms



Image from eSchoolToday: Revision notes, Soil


Soil Carbon

- Soil has about 1500 Pg of carbon worldwide
- As soil is heated, it releases carbon
 - Not all soil is the same
 - Plants and microbes affect carbon loss rate



Soil Nitrogen

- Soil has about 133–140 Pg of nitrogen worldwide
- As soil is heated, it can speed up mineralization (decay)
 - Could be offset by other parts of the nitrogen cycle speeding up



	Previous studies	Our study
Time	Long term 6 months-7 years	Short term 10 days
Degrees heating	Gradual, similar to greenhouse effect 1-5°C increase	Extreme, similar to heat wave 30-60°C increase
Methods	Overhead heating lamp Greenhouses Buried cables	Overhead infra-red heating lamp (sunlight-like source)
Results	Nitrogen mineralization (N loss)* Increased CO ₂ flux (C loss)* Changing plant/bacteria responses	?


*Dependent on hydrology

Will extreme short-term heating show similar soil carbon and nitrogen losses as long-term experiments?

Hypothesis

- Carbon and nitrogen loss from soil
- More loss if watered
 - Due to microbial activity and respiration

Sample Expected Results



Site Background

- Oak Woodlands cover 10 million acres of California
 - Average carbon content: 4.1% (min 3.0% max 6.5%)
 - Average nitrogen content: 0.28% (min 0.15% max 0.45%).



- Hastings Natural History Reservation (HNHR)
 - Biological field station of Museum of Vertebrate Zoology at UC Berkeley since 1937
 - Coast Live Oak (*Quercus agrifolia*) and grassland




Image from "North American Oak Woodland" from University of Arizona, College of Agriculture and Life Sciences.

Methods: Gathering Samples



Remove plant cover



Cut ~30x3x5cm soil sample



Transfer to labeled bag

Methods: Processing Soil



Sift through sieve



Remove large roots & seeds




Separate into thirds
(~150g-200g each)

Methods: Heating Soil

Heat lamps, 6 hours a day, 10 days total




10 samples



10 samples

35mL water
every 5 days



10 samples



Half the samples were heated at high (direct) heat; half were heated at low heat.

After treatment, samples were mailed to OSU's Central Analytical Laboratory (Crop and Soil Science Dept) for analysis




What I expect to find...

Hypothesis


- Carbon and nitrogen loss from soil
- More loss if watered
 - Due to microbial activity and respiration

Sample Expected Results



Soil Carbon


	Dry Treatment	Wet Treatment
Soil C increases	7	7
Soil C decreases	2	3
Soil C stays the same	1	0



- No clear trend
- Changes are very small
- Average Carbon = $3.78\% \pm 1.02$

Soil Nitrogen


	Dry Treatment	Wet Treatment
Soil N increases	5	7
Soil N decreases	5	3
Soil N stays the same	0	0



- No clear trend
- Changes are very small
- Average Nitrogen = $0.29\% \pm 0.08$

Conclusions

Extreme short-term warming seems to be ineffective in reducing carbon and nitrogen in Oak Woodlands soil.



Further Questions

Extreme short-term warming seems to be ineffective in reducing carbon and nitrogen in Oak Woodlands soil.

Should short-term warming show the same results as long term?

- Not necessarily

What if there is CO_2 being lost, but it's not enough to show?

- Previous studies: around 0.001 to $0.006 \text{ mgC cm}^{-2} \text{ hr}^{-1}$ net loss comparing control plots and heated plots
- Our study: around $19610 \text{ mgC cm}^{-2}$ in the soil sample, 60 hours of heating
 - Result: only 0.36 mgC cm^{-2} loss

Classroom Research



- Thomas Jefferson High School IB Environmental Science students
- Long-term extreme soil heating study
- Student-run
- Ongoing through the school year

Sources Cited

&



Acknowledgements

- Bai, E.; Li, S.; Xu, W.; Li, W.; Dai, W. & Jiang, P. (2013). *A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics*. New Phytologist **199**:441-451. DOI: 10.1111/nph.12
- Batjes, N.H. (2014). *Total carbon and nitrogen in the soils of the world*. European Journal of Soil Science. <https://doi.org/10.1111/ejss.12114> 2
- *Carbon Pools and Fluxes*. c2008. Durham (NH): University of New Hampshire; [accessed 2018 August 1]. <http://globecarboncycle.unh.edu/CarbonPoolsFluxes.shtml>
- Dahlgren, R.A.; Horwath, W.R.; Tate, K.W.; Camping, T.J. (2003). *Blue oak enhance soil quality in California oak woodlands*. Hilgardia 57(2):42-47. DOI: 10.3733/ca.v057n02p42
- Lu X, Fan J, Yan Y, Wang X (2013) Responses of Soil CO₂ Fluxes to Short-Term Experimental Warming in Alpine Steppe Ecosystem, Northern Tibet. PLoS ONE 8(3): e59054. doi:10.1371/journal.pone.0059054
- Melillo, J.M.; Frey, S. D.; DeAngelis, K.M.; Werner, W.J.; Bernard, M.J.; Bowles, F.P.; Pold, G.; Knorr, M.A. & Grandy, A.S. (2017). *Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world*. Science **358** (6359), 101-105. DOI: 10.1126/science.aan2874
- Melillo, J.M.; Steudler, P.A.; Aber, J.D.; Newkirk, K.; Lux, H.; Bowles, F.P.; Catricala, C.; Magill, A.; Ahrens, T.; Morrisseau, S. (2002). *Soil Warming and Carbon-Cycle Feedbacks to the Climate System*. Science **298** (5601), 2173-2176. DOI: 10.1126/science.1074153
- *North American Oak Woodland*. c2018. Arizona: University of Arizona (College of Agriculture and Life Sciences); [accessed 2018 July 29]. <https://wrangle.org/ecotype/north-american-oak-woodlands>
- Rice, C.W. (2002). *"Storing carbon in soil: Why and how?"*. Geotimes. **47** (1): 14-17. Retrieved 7 August, 2018.
- Scharlemann, J.; Tanner, E.; Hiederer, R.; & Kapos, V. (2014) *Global soil carbon: understanding and managing the largest terrestrial carbon pool*. Carbon Management, 5:1, 81-91, DOI: 10.4155/cmt.13.77
- Silver, W.L.; Ryals, R.; Evine, V. (2010). *Soil Carbon Pools in California's Annual Grassland Ecosystem*. Rangeland Ecology Management **63**: 128-136. DOI: 10.2111/REM-D-09-0010
- Waldrop, M.P.; Firestone, M.K. (2004). *Microbial community utilization of recalcitrant and simple carbon compounds: impact of oak-woodland plant communities*. Ecosystem Ecology **138**: 275-284. DOI 10.1007/s00442-003-1419-9



Thanks to...

- My entire STAR family
- My mentor, Chris
- My Student Scientists from TJHS!
 - Rachel
 - Kyle
 - Rebecca
 - Dennis

The 2018 STEM Teacher and Researcher Program and this project have been made possible through support from Chevron (www.chevron.com), the National Marine Sanctuary Foundation (www.marinesanctuary.org), the National Science Foundation through the Robert Noyce Program under Grant #1836335 and 1340110, the California State University Office of the Chancellor, and California Polytechnic State University in partnership with NASA Ames Research Center. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the funders.



Carbon	Average	Stdev	Coef of Var
Control	3.7	1.06	28.59
Low Dry	3.8	1.08	28.47
High Dry	3.7	0.99	26.56
Low Wet	3.9	1.36	35.43
High Wet	3.9	0.98	25.48



Nitrogen	Average	Stdev	Coef of Var
Control	0.29	0.08	27.51
Low Dry	0.28	0.08	28.42
High Dry	0.28	0.08	26.89
Low Wet	0.29	0.09	30.42
High Wet	0.30	0.09	29.15